

Schwinger-Dyson BRST-Symmetry and the Equivalence of Hamiltonian and Lagrangian Quantisation

FRANK DE JONGHE*

Instituut voor Theoretische Fysica, K.U.Leuven, Belgium
and
CERN – Geneva

February 1, 2008

Abstract

Implementing the requirement that a field theory be invariant under Schwinger-Dyson BRST symmetry in the Hamiltonian formalism, we show the equivalence between Hamiltonian and Lagrangian BRST-formalism at the path integral level. The Lagrangian quantum master equation is derived as a direct consequence of the Fradkin-Vilkovisky theorem in Hamiltonian BRST quantisation.

KUL-TF-93/08
CERN-TH-6823/93
March 1993

*Aspirant of the N.F.W.O., Belgium.

There are two approaches available for the quantisation of gauge theories, both rooted in the concept of BRST symmetry. First, a Hamiltonian method was developed (below denoted by BFV) [1], with as key features a dynamical treatment of the Lagrange multipliers imposing the constraints, the introduction of a canonically conjugate pair of a ghost and an antighost for every first-class constraint, the existence of a nilpotent fermionic BRST-generator, and the existence of a BRST invariant Hamiltonian. Using these ingredients one can construct a path integral which is independent of the gauge fixing, as guaranteed by the celebrated Fradkin-Vilkovisky theorem [1]. Guided by but not as a logical consequence of this Hamiltonian formalism a Lagrangian formalism was developed (below denoted by BV)[2]. Here, the most important property is the doubling of the fields, by associating with every field ϕ^a an antifield ϕ_a^* of opposite statistics. Solving the so-called master equation gives an extended action, depending on both the fields and the anti-fields. The gauge fixing is done by replacing the antifield ϕ_a^* by $\frac{\delta\Psi(\phi)}{\delta\phi^a}$, where Ψ is a fermionic function of the fields, called the gauge fermion [1, 2]. In this paper we will prove the equivalence of both methods for quantisation.

A large effort has already been devoted over the years to precisely this problem. In a series of papers [3], Henneaux et al. proved the following. Given any Lagrangian, one can try to find its gauge symmetries directly, and treat them in the scheme of BV. On the other hand, one could look for the gauge symmetries using the Hamiltonian methods of Dirac. The canonical action so obtained can again be treated in the BV scheme. Both methods were proven to lead to the same path integral. The most remarkable feature of this approach is that in the Hamiltonian method, anti-fields have to be introduced for the momenta. The Lagrangian analog of the Hamiltonian BRST-charge has also been constructed in [4].

The procedure we will follow here, is closer to that of [5]. There it was shown that starting from a complete Hamiltonian analysis following BFV, it is possible to obtain a Lagrangian action satisfying the quantum master equation of BV, after integrating out the momenta. However, the anti-fields have to be introduced in the Hamiltonian path integral in a rather ad hoc way as sources for the BRST-transformation. Also, the Lagrangian master equation requires a separate proof, where one would expect it to follow naturally from the Fradkin-Vilkovisky theorem, as they both have as consequence that the generating function does not depend on the choice gauge. We will clarify these two points below.

Recently, it was shown [7] that demanding a Lagrangian field theory to be invariant under Schwinger-Dyson BRST symmetry [6] in addition to usual BRST symmetries of internal gauge invariances is a natural way of introducing the anti-fields of the Batalin-Vilkovisky Lagrangian formalism. It turns out that they are the anti-ghosts of the Schwinger-Dyson BRST symmetry. The derivation is done most simply in terms of a certain Lagrangian collective field formalism, described in detail in ref. [6], but can also be based directly on the Schwinger-Dyson BRST symmetry itself. However, it is less obvious how to implement the latter approach in the Hamiltonian formalism.

Our strategy to prove the equivalence of the Hamiltonian BFV formalism and the Lagrangian BV formalism will then be to consider a Hamiltonian system with known extended phase space, BRST-charge and BRST-invariant Hamiltonian. These are all

constructed following the prescriptions of the BFV formalism [8]. Then we implement the Schwinger-Dyson shift symmetry in a way consistent with the original symmetries. Extending further the extended phase space to include the ghost-antighost pairs for this shift symmetry, we introduce the anti-fields. Gauge fixing the shift-symmetry and the original gauge-symmetries allows one to construct the path-integral. After integrating out all momenta, except the ones for the Lagrange multipliers of the Hamiltonian system, one ends up with an action satisfying the BV quantum master equation. Anti-fields indeed act as sources for BRST-transformations and the left-over momenta become the Nakanishi-Lautrup fields.

We will first recapitulate how Schwinger-Dyson shift symmetries are to be described in the Hamiltonian formalism [6]. To that purpose we first consider a Lagrangian depending on fields and their time-derivatives -denoted by a dot- which has no internal gauge symmetries : $L(t) = L(\phi^a, \dot{\phi}^a)$. Introducing collective fields amounts here to considering $L(\phi^a - \varphi^a, \dot{\phi}^a - \dot{\varphi}^a)$. Denoting the momenta conjugate to ϕ^a by π_a and to φ^a by ϖ_a , the first class constraints are

$$\chi_a = \pi_a + \varpi_a = 0 \quad (1)$$

From this, it trivially follows that the structure constants C_{bc}^a vanish, $[\chi_b, \chi_c] = 0$, as $[\pi_a, \phi^b] = [\varpi_a, \varphi^b] = -\delta_a^b$ are the only non-vanishing Poisson brackets. These constraints also have a vanishing Poisson bracket with the Hamiltonian, as the latter also only depends on the difference $\phi^a - \varphi^a$. This is equivalent to saying that the structure constants V_a^b vanish. After associating one Lagrange multiplier λ^a and its canonical momentum $\pi_{\lambda a}$ ($[\pi_{\lambda a}, \lambda^b] = -\delta_a^b$) with every first class constraint χ_a the complete set of constraints $G_a = (\pi_{\lambda a}, \chi_a)$ still has vanishing structure constants. We now introduce the following ghost and antighost fields to construct the extended phase space [8]

$$\begin{aligned} \pi_{\lambda a} &= 0 \quad \begin{array}{l} gh_{\leftrightarrow}^{\sharp=1} \\ gh_{\leftrightarrow}^{\sharp=-1} \end{array} \quad \begin{array}{l} -i^{a+1}\mathcal{P}^a \\ i^{a+1}\phi_a^* \end{array} \\ \chi_a &= 0 \quad \begin{array}{l} gh_{\leftrightarrow}^{\sharp=1} \\ gh_{\leftrightarrow}^{\sharp=-1} \end{array} \quad \begin{array}{l} c^a \\ \bar{\mathcal{P}}_a \end{array} \end{aligned} \quad (2)$$

with the only non-vanishing brackets $[\bar{\mathcal{P}}_a, c^b] = [\mathcal{P}^b, \phi_a^*] = -\delta_a^b$. The ghostfields (those of $gh_{\sharp} = 1$) are collectively denoted by η^a , the anti ghosts ($gh_{\sharp} = -1$) by θ_a , including the numerical factors i in (2). Using this, one can straightforwardly construct the BRST generator of the shift symmetries

$$\Omega_s = -i^{a+1}\mathcal{P}^a\pi_{\lambda a} + c^a(\pi_a + \varpi_a) \quad (3)$$

Gauge fixing the collective field to zero can then be done by taking as gauge-fermion

$$\psi = \bar{\mathcal{P}}_a\lambda^a + \frac{i^{a+1}}{\beta}\phi_a^*\varphi^a \quad (4)$$

where β is an arbitrary parameter. We will take the $\beta \rightarrow 0$ limit later. The Poisson bracket with the BRST-charge then gives

$$[\psi, \Omega_s] = i^{a+1}\mathcal{P}^a\bar{\mathcal{P}}_a - \frac{1}{\beta}\varphi^a\pi_{\lambda a} - \lambda^a(\pi_a + \varpi_a) + \frac{i^{a+1}(-1)^a}{\beta}\phi_a^*c^a \quad (5)$$

The action to be exponentiated in the path-integral is then according to the standard BfV formalism given by

$$S = \dot{\phi}^a \pi_a + \dot{\varphi}^a \varpi_a + \dot{\lambda}^a \pi_{\lambda a} + \dot{\eta}^a \theta_a - H(\pi, \varpi, \phi - \varphi) + [\psi, \Omega_s] \quad (6)$$

We now redefine

$$\begin{aligned} \frac{1}{\beta} \pi_{\lambda a} &\rightarrow \pi_{\lambda a} \\ \frac{i^{a+1}}{\beta} \phi_a^* &\rightarrow \phi_a^* \\ (-1)^a c^a &\rightarrow c^a \end{aligned} \quad (7)$$

At this moment we take the limit $\beta \rightarrow 0$, which makes two terms of the symplectic part of S disappear. After that, the integral over the ghostmomenta $\mathcal{P}^a$ and $\bar{\mathcal{P}}_a$ becomes trivial, leading to

$$S = \dot{\phi}^a \pi_a + \dot{\varphi}^a \varpi_a - H(\pi, \varpi, \phi - \varphi) - \varphi^a \pi_{\lambda a} - \lambda^a (\pi_a + \varpi_a) + \phi_a^* c^a \quad (8)$$

By integrating over the Lagrange multipliers and their momenta, the path integral becomes

$$\int [d\phi^a][dc^a][d\phi_a^*] e^{\frac{i}{\hbar}[S + \phi_a^* c^a]} \quad (9)$$

where

$$e^{\frac{i}{\hbar}S} = \int [d\pi_a] \exp \left[\frac{i}{\hbar} \left(\int dt \dot{\phi}^a \pi_a - H(\pi, -\varpi, \phi) \right) \right] \quad (10)$$

This parallels the result of [7] for the case of no gauge-symmetries. The action is nothing but the original action, plus a term where the anti-fields multiply the ghosts c^a of Alfaro and Damgaard [7]. Notice also that the collective field has been eliminated completely. Recently, it was shown [9] that this is not possible when one tries to construct a path integral which is invariant under both BRST and anti-BRST transformations. The collective field then is needed as a source for mixed transformations [10].

We want to add another comment, which has some bearing on the case where there are originally gauge-symmetries present. The Hamiltonian introduced in (6) has a rather arbitrary dependence on the momenta. For instance, if we had started from a Lagrangian quadratic in the velocities, $L(t) = \frac{1}{2}\dot{\phi}^2 + \tilde{L}(\phi)$ then an arbitrary parameter could be present in the Hamiltonian

$$H = \frac{1}{2}\mu\pi^2 + \frac{1}{2}(1 - \mu)\varpi^2 - \tilde{L}(\phi - \varphi) \quad (11)$$

Of course, the μ -dependence disappears when imposing the constraint. When there are gauge symmetries present, a specific choice of μ is needed to make contact with BV.

Now we turn to the case of a Hamiltonian with gauge symmetries. We start from an extended phase space, whose coordinates $(\phi^a, c^a, \bar{c}_a, \lambda^a$ for the case of first class irreducible theories) we collectively denote by $Y^{\bar{a}}$, the conjugate momenta $(\pi_a, \bar{\mathcal{P}}_a, \mathcal{P}^a, \pi_{\lambda a})$ by $\Pi_{\bar{a}}$. The fundamental Poisson bracket is $[\Pi_{\bar{a}}, Y^{\bar{b}}] = -\delta_{\bar{a}}^{\bar{b}}$. The fact that the symplectic form

might be more complicated after the elimination of second-class constraints, can easily be incorporated. We start from a known BRST-charge $\Omega_0(\Pi_{\bar{a}}, Y^{\bar{a}})$ and BRST-invariant Hamiltonian $H_0(\Pi_{\bar{a}}, Y^{\bar{a}})$ that satisfy the standard conditions $[\Omega_0, \Omega_0] = [H_0, \Omega_0] = 0$.

We now introduce for every field $Y^{\bar{a}}$ a collective field $\mathcal{Y}^{\bar{a}}$ which has conjugate momenta $\Upsilon_{\bar{a}}$, again with fundamental Poisson bracket $[\Upsilon_{\bar{a}}, \mathcal{Y}^{\bar{b}}] = -\delta_{\bar{a}}^{\bar{b}}$. It will be useful to associate with every phase space function $F(\Pi_{\bar{a}}, Y^{\bar{a}})$ a function $\tilde{F} = F(-\Upsilon_{\bar{a}}, Y^{\bar{a}} - \mathcal{Y}^{\bar{a}})$. It is then straightforward to show that

$$[\tilde{F}, \tilde{G}] = [\widetilde{F}, \widetilde{G}] \quad (12)$$

As the constraints of the shift symmetry are still given by $\chi_{\bar{a}} = \Pi_{\bar{a}} + \Upsilon_{\bar{a}}$, it also holds that

$$[\tilde{F}, \chi_{\bar{a}}] = 0 \quad (13)$$

In order to construct the correct Hamiltonian for which both the original gauge symmetries and the shift symmetries are present, we look for

$$\Omega = \widetilde{\Omega}_0 + \Omega_s \quad (14)$$

$$H = \widetilde{H}_0 + \Delta H \quad (15)$$

and demand that $[\Omega, \Omega] = [H, \Omega] = 0$.

Constructing the extended phase space as in the case of no internal symmetries

$$\begin{aligned} \pi_{\lambda_{\bar{a}}} &= 0 & \begin{array}{l} gh_{\leftrightarrow}^{\sharp=1} \\ gh_{\leftrightarrow}^{\sharp=-1} \end{array} & \begin{array}{l} -i^{\bar{a}+1} \mathcal{P}^{\bar{a}} \\ i^{\bar{a}+1} Y_{\bar{a}}^* \end{array} \\ \chi_a &= 0 & \begin{array}{l} gh_{\leftrightarrow}^{\sharp=1} \\ gh_{\leftrightarrow}^{\sharp=-1} \end{array} & \begin{array}{l} c^{\bar{a}} \\ \bar{\mathcal{P}}_{\bar{a}} \end{array} \end{aligned} \quad (16)$$

the total BRST-charge is obtained by taking for Ω_s the same thing as in the case without gauge symmetries.

$$\Omega = \widetilde{\Omega}_0 - i^{\bar{a}+1} \mathcal{P}^{\bar{a}} \pi_{\lambda_{\bar{a}}} + c^{\bar{a}} (\Pi_{\bar{a}} + \Upsilon_{\bar{a}}) \quad (17)$$

That this quantity satisfies $[\Omega, \Omega] = 0$ follows immediately from (12) and (13). From these it is also clear that $\Delta H = 0$, i.e. $H = \widetilde{H}_0$. In fact, these two properties simply reflect the vanishing of the structure constants associated with the Poisson brackets of the original constraints and the original Hamiltonian with the constraints of the shift symmetry, when the former are evaluated in $(-\Upsilon_{\bar{a}}, Y^{\bar{a}} - \mathcal{Y}^{\bar{a}})$.

One might have wondered why the $\tilde{F}$ operation also involves a substitution of the momenta by minus the momenta of the collective field. This becomes clear when evaluating how the fields and the collective fields transform under the new BRST-charge. The fields only transform under the shift symmetry

$$[Y^{\bar{a}}, \Omega] = (-1)^{\bar{a}} c^{\bar{a}} \quad (18)$$

while the original gauge transformations have shifted to the collective fields

$$[\mathcal{Y}^{\bar{a}}, \Omega] = (-1)^{\bar{a}} c^{\bar{a}} + [\mathcal{Y}^{\bar{a}}, \widetilde{\Omega}_0] \quad (19)$$

The original transformation can be shuffled between the fields and their collective partners, which is related to the arbitrariness in the momentum dependence of the Hamiltonian, as exemplified in (11). It is precisely by our momentum substitution rule that they end up entirely in the collective field transformation. It was shown in [7] that this corresponds to the BV boundary condition that the solution of the classical master equation starts with terms linear in the anti-fields, i.e. source terms for the BRST formations. Other choices lead to equally well-defined path integrals but they do not necessarily agree with the BV boundary conditions.

For the gauge-fixing of the collective field we use again (4)

$$\psi = \bar{\mathcal{P}}_{\bar{a}} \lambda^{\bar{a}} + \frac{i^{\bar{a}+1}}{\beta} Y_{\bar{a}}^* \mathcal{Y}^{\bar{a}} \quad (20)$$

leading to

$$[\psi, \Omega] = i^{\bar{a}+1} \mathcal{P}^{\bar{a}} \bar{\mathcal{P}}_{\bar{a}} - \frac{1}{\beta} \mathcal{Y}^{\bar{a}} \pi_{\lambda \bar{a}} - \lambda^{\bar{a}} (\Pi_{\bar{a}} + \Upsilon_{\bar{a}}) + \frac{i^{\bar{a}+1} (-1)^{\bar{a}}}{\beta} Y_{\bar{a}}^* c^{\bar{a}} + \frac{i^{\bar{a}+1}}{\beta} Y_{\bar{a}}^* [\mathcal{Y}^{\bar{a}}, \widetilde{\Omega}_0] \quad (21)$$

where the last term is the only, but important, difference from the case of no gauge symmetries. We can rewrite its Poisson bracket as

$$[\mathcal{Y}^{\bar{a}}, \widetilde{\Omega}_0] = [\widetilde{Y}^{\bar{a}}, \widetilde{\Omega}_0] = [Y^{\bar{a}}, \widetilde{\Omega}_0] \quad (22)$$

because $\widetilde{\Omega}_0$ does not depend on the momentum conjugate to $Y^{\bar{a}}$.

The gauge fixing of the original symmetry can be done by taking a fermion $\Psi(\Pi_{\bar{a}}, Y^{\bar{a}})$ and adding

$$[\widetilde{\Psi}, \Omega] = [\widetilde{\Psi}, \widetilde{\Omega}_0] \quad (23)$$

Doing the same field-redefinitions as in (7), of course with a replacement of the indices a by $\bar{a}$, we get the standard BFM action

$$\begin{aligned} S = & \dot{Y}^{\bar{a}} \Pi_{\bar{a}} + \dot{\mathcal{Y}}^{\bar{a}} \Upsilon_{\bar{a}} + \beta \dot{\lambda}^{\bar{a}} \pi_{\lambda \bar{a}} - \beta i^{\bar{a}+1} \dot{\mathcal{P}}^{\bar{a}} Y_{\bar{a}}^* + (-1)^{\bar{a}} \dot{c}^{\bar{a}} \bar{\mathcal{P}}_{\bar{a}} \\ & - \widetilde{H}_0 + i^{\bar{a}+1} \mathcal{P}^{\bar{a}} \bar{\mathcal{P}}_{\bar{a}} - \mathcal{Y}^{\bar{a}} \pi_{\lambda \bar{a}} - \lambda^{\bar{a}} (\Pi_{\bar{a}} + \Upsilon_{\bar{a}}) \\ & + Y_{\bar{a}}^* c^{\bar{a}} + Y_{\bar{a}}^* [Y^{\bar{a}}, \widetilde{\Omega}_0] + [\widetilde{\Psi}, \widetilde{\Omega}_0] \end{aligned} \quad (24)$$

After taking the limit $\beta \rightarrow 0$, the ghost-momenta $\mathcal{P}$ and $\bar{\mathcal{P}}$ can be integrated out trivially. Doing then the integral over $\pi_{\lambda \bar{a}}$ gives a delta-function $\delta(\mathcal{Y}^{\bar{a}})$, while integrating out the Lagrange multiplier $\lambda^{\bar{a}}$ itself leads to $\delta(\Pi_{\bar{a}} + \Upsilon_{\bar{a}})$. It is trivial to see that imposing these delta-function constraints allows us to drop the tildes, so that after integrating over the collective field and its momentum we get

$$S = \dot{Y}^{\bar{a}} \Pi_{\bar{a}} - H_0 + Y_{\bar{a}}^* [Y^{\bar{a}}, \Omega_0] + Y_{\bar{a}}^* c^{\bar{a}} + [\Psi, \Omega_0] \quad (25)$$

We shall now show that the path integral obtained from this action, is equal to the one obtained by doing Lagrangian BV analysis. We make the most popular choice for the

gauge fermion ¹

$$\Psi = \Psi_0(Y^{\bar{a}}, \pi_{\lambda a}) + \bar{\mathcal{P}}_a \lambda^a \quad (26)$$

Furthermore, we know the form of Ω_0

$$\Omega_0 = -i^{a+1} \mathcal{P}^a \pi_{\lambda a} + \Omega_{min} \quad (27)$$

where Ω_{min} does not depend on the Lagrange multiplier λ^a nor its momentum $\pi_{\lambda a}$. This allows us to write

$$[\Psi, \Omega_0] = \frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}} [Y^{\bar{a}}, \Omega_0] + i^{a+1} \mathcal{P}^a \bar{\mathcal{P}}_a - (-1)^a \frac{\delta^r \Omega_{min}}{\delta c^a} \lambda^a \quad (28)$$

It is now convenient to define $\hat{S}$ by

$$S = \hat{S}(Y^{\bar{a}}, Y_{\bar{a}}^*, \pi_{\lambda a}, \mathcal{P}^a, \bar{\mathcal{P}}_a, \pi_\alpha) + Y_{\bar{a}}^* c^{\bar{a}} + \frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}} \frac{\delta^l \hat{S}}{\delta Y_{\bar{a}}^*} \quad (29)$$

so that the complete path integral, obtained by Hamiltonian methods and by integrating a part of the extended phase space associated with the shift symmetry, is given by

$$\int [dY^{\bar{a}}][d\pi_{\lambda a}][dY_{\bar{a}}^*][dc^{\bar{a}}][d\mathcal{P}^a][d\bar{\mathcal{P}}_a][d\pi_\alpha] \exp \left[\frac{i}{\hbar} Y_{\bar{a}}^* c^{\bar{a}} \right] \hat{U} \left(\exp \left[\frac{i}{\hbar} \hat{S} \right] \right) \quad (30)$$

where the operator $\hat{U}$ is defined by

$$\hat{U} = \exp \left[\frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}} \frac{\delta^l}{\delta Y_{\bar{a}}^*} \right] \quad (31)$$

We now define $\mathcal{S}$ by

$$e^{\frac{i}{\hbar} \mathcal{S}(Y^{\bar{a}}, Y_{\bar{a}}^*)} = \int [d\mathcal{P}^a][d\bar{\mathcal{P}}_a][d\pi_\alpha] e^{\frac{i}{\hbar} \hat{S}} \quad (32)$$

Since the operator $\hat{U}$ commutes with the integrations done to define $\mathcal{S}$, we can rewrite the partition function as

$$\int [dY^{\bar{a}}][d\pi_{\lambda a}][dY_{\bar{a}}^*][dc^{\bar{a}}] \exp \left[\frac{i}{\hbar} \left(Y_{\bar{a}}^* c^{\bar{a}} + \mathcal{S}(Y^{\bar{a}}, Y_{\bar{a}}^* + \frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}}) \right) \right] \quad (33)$$

where we used the well-known relation $\exp \left[a(y) \frac{\delta}{\delta x} \right] f(x) = f(x + a(y))$. Integrating over the ghost $c^{\bar{a}}$ leads to a $\delta(Y_{\bar{a}}^*)$, removing the anti-fields. We thus recover the gauge fixing procedure of [2], which is to start from an extended action, function of fields and anti-fields, and to replace the latter by a derivative of some gauge fermion w.r.t. its conjugate field.

¹The Lagrange multipliers, ghosts and anti ghosts that appear below, must not be confused with the same quantities for the shift symmetries which have been integrated over above. The former have indices a , while the latter had $\bar{a}$. The point is that to proceed further we have for the first time to specify in detail what fields are contained in $Y^{\bar{a}}$ and $\Pi_{\bar{a}}$.

In contrast to [5], we do not integrate over the Lagrange multipliers, but include them together with their momenta in the set of degrees of freedom of the obtained Lagrangian system. This is done because they are usually needed to get a covariant Lagrangian in the end, as the example of Yang-Mills theories teaches us. There the Lagrange multipliers are nothing but the fourth component of the Lorentz-vector A_μ^a .

It is now trivial to derive the Lagrangian quantum master equation, satisfied by $\mathcal{S}$. We use the Fradkin-Vilkovisky theorem, which states that we are allowed to change Ψ_0 to $\Psi_0 + \Delta\Psi$ without changing the partition function. For an infinitesimal change in the gauge fermion we thus get

$$\int [dY^{\bar{a}}][dY_{\bar{a}}^*][dc^{\bar{a}}][d\pi_{\lambda_a}] e^{+\frac{i}{\hbar} Y_{\bar{a}}^* c^{\bar{a}}} \frac{\delta^r \Delta\Psi}{\delta Y^{\bar{a}}} \frac{\delta^l}{\delta Y_{\bar{a}}^*} e^{\frac{i}{\hbar} \hat{\mathcal{S}}} = 0 \quad (34)$$

for any choice for $\Delta\Psi$. We denoted $\mathcal{S}(Y^{\bar{a}}, Y_{\bar{a}}^* + \frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}}) = \hat{\mathcal{S}}$. Because of the arbitrariness of $\Delta\Psi$, this leads after a partial integration to the well-known Quantum Master Equation [2]

$$0 = (-1)^{\bar{a}} \frac{\delta^l}{\delta Y^{\bar{a}}} \frac{\delta^l}{\delta Y_{\bar{a}}^*} e^{\frac{i}{\hbar} \hat{\mathcal{S}}} = \Delta e^{\frac{i}{\hbar} \hat{\mathcal{S}}} \quad (35)$$

which is equivalent to the more familiar form

$$(\hat{\mathcal{S}}, \hat{\mathcal{S}}) = 2i\hbar \Delta \hat{\mathcal{S}} \quad (36)$$

The so-called anti-bracket is defined as

$$(F, G) = \frac{\delta^r F}{\delta Y^{\bar{a}}} \frac{\delta^l G}{\delta Y_{\bar{a}}^*} - \frac{\delta^r F}{\delta Y_{\bar{a}}^*} \frac{\delta^l G}{\delta Y^{\bar{a}}} \quad (37)$$

Instead of gauge fixing the original symmetries as done in (23), we could also just have added

$$[\Psi_0(Y^{\bar{a}}, \pi_{\lambda_a}) + \widetilde{\mathcal{P}_a} \lambda^a, \Omega] = -\frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}} c^{\bar{a}} + [\bar{\mathcal{P}}_a \lambda^a, \Omega_0] \quad (38)$$

where we have done the redefinition (7), and used that integrating over the collective field and its momentum makes the tildes disappear. If we now integrate over the $c^{\bar{a}}$ ghost, we get as gauge fixing condition

$$\delta(Y_{\bar{a}}^* - \frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}}) \quad (39)$$

Redefining

$$Y_{\bar{a}}^* - \frac{\delta^r \Psi_0}{\delta Y^{\bar{a}}} \rightarrow Y_{\bar{a}}^* \quad (40)$$

one easily sees that this is exactly the same partition function as before.

Let us now try to get a better understanding of $\mathcal{S}(Y, Y^*)$ defined in (32). Following the usual procedure for solving the quantum master equation [2], we expand it in a series in $\hbar$

$$\mathcal{S} = S_0 + \sum_{i=1}^{+\infty} \hbar^i M_i \quad (41)$$

We will only discuss S_0 , which satisfies the classical master equation $(S_0, S_0) = 0$. It can be calculated by applying the saddle point approximation to (32). Solving the field equations

$$\frac{\delta \hat{S}}{\delta \mathcal{P}^a} = \frac{\delta \hat{S}}{\delta \bar{\mathcal{P}}_a} = \frac{\delta \hat{S}}{\delta \pi_\alpha} = 0 \quad (42)$$

gives us functions $\mathcal{P}^a(Y^{\bar{a}}, Y_{\bar{a}}^*, \pi_{\lambda a})$ and $\bar{\mathcal{P}}_a(Y^{\bar{a}}, Y_{\bar{a}}^*, \pi_{\lambda a})$ and $\pi_\alpha(Y^{\bar{a}}, Y_{\bar{a}}^*, \pi_{\lambda a})$. Then we have

$$S_0(Y, Y^*, \pi_{\lambda a}) = \hat{S}|_\Sigma \quad (43)$$

where the notation $|_\Sigma$ means that we plugged in the solutions of the extremum equations. If we now define

$$(Y^{\bar{a}}, S_0)|_{Y^*=0} = \mathcal{R}_{\bar{a}}^a c^a \quad (44)$$

then it is trivial to show that

$$\mathcal{R}_{\bar{a}}^a c^a = [Y^{\bar{a}}, \Omega_0]|_{\Sigma, Y^*=0} \quad (45)$$

so that indeed the part of S_0 linear in the anti-fields acts as a source term for the BRST-transformations.

This finishes our proof of the equivalence of the Hamiltonian BFV and the Lagrangian BV formalism. Our guiding principle was that the Schwinger-Dyson shift symmetry allows for a natural introduction of anti-fields [7]. Using the prescriptions of the BFV scheme to implement the Schwinger-Dyson BRST symmetry, we see that the presence of the anti-fields need not be restricted to Lagrangian BV. However, the most obvious manipulations lead straightforward to an interpretation like that of the well-known Lagrangian scheme of BV. The Lagrangian action we get in the end, trivially satisfies the BV quantum master equation, as a result of the Fradkin-Vilkovisky theorem. We thus have linked the two principles which assure that the Hamiltonian and Lagrangian method can be used for quantising gauge theories at all, namely, that the partition functions constructed following their prescription, is independent of the chosen gauge fixing.

It has been known for some time, that anomalies occur in the BV scheme when one can not construct a local M_1 when trying to solve the master equation at $\mathcal{O}(\hbar)$ [11]. To our knowledge, there have been no studies on how anomalies would arise in the regularised BFV path integrals. It would be interesting to see how the momentum integrations play a role in relating both schemes on that account.

Although both our starting point, the Hamiltonian BFV, and our end point, the Lagrangian BV, are capable of handling theories with open algebras, it is not clear how the collective field formalism should be modified in that case [7]. Clearly, this is one of the major points to be clarified in the future.

Acknowledgment

It is a real pleasure to thank P.H. Damgaard for many vivid discussions, and for making my stay at CERN both enjoyable and mentally profitable. I also want to thank W. Troost and A. Van Proeyen for sharing their insights on BV.

References

- [1] E.S. Fradkin and G.A. Vilkovisky, Phys.Lett. **B55** (1975) 224
I.A. Batalin and G.A. Vilkovisky, Phys.Lett. **B69** (1977) 309
E.S. Fradkin and T.E. Fradkina, Phys.Lett. **B72** (1978) 343
I.A. Batalin and E.S. Fradkin, Phys.Lett. **B122** (1983) 157
- [2] I.A. Batalin and G.A. Vilkovisky, Phys. Lett. **102B** (1981) 27; Phys. Rev. **D28** (1983) 2567 [E: **D30** (1984) 508]; Nucl. Phys. **B234** (1984) 106; J. Math. Phys. **26** (1985) 172.
- [3] J.Fisch and M. Henneaux, Phys.Lett. **B226** (1989) 80
M. Henneaux, Phys.Lett. **238** (1990) 299
A. Dresse, Ph. Gregoire, M. Henneaux, Phys.Lett. **245** (1990) 192
A. Dresse, J. Fisch, P. Gregoire and M. Henneaux Nucl.Phys. **B354** (1991) 191
- [4] C.Battle, J.Gomis, J.Paris and J.Roca, Phys.Lett. **B224** (1989) 288 and Nucl.Phys. **239** (1990) 139
W.Siegel, Int.J.Mod.Phys. A Vol **4** (1989) 3951
G.V. Grigoryan, R.P. Grigoryan and I.V. Tyutin, Nucl.Phys **B379** (1992) 304
- [5] G.V. Grigoryan, R.P. Grigoryan and I.V. Tyutin, Sov.J.Nucl.Phys. **53** (1991) 1058
- [6] J. Alfaro and P.H. Damgaard, Phys. Lett. **B222** (1989)425 and Ann. Phys.(N.Y.) **202** (1990) 398.
J. Alfaro, P.H. Damgaard, J. Latorre and D. Montano, Phys. Lett. **B233** (1989) 153.
- [7] J. Alfaro and P.H. Damgaard, preprint CERN–TH-6788/93, hep-th 9301103.
- [8] M. Henneaux, Phys.Rep. **126** (1985) 1
J. Govaerts, Hamiltonian Quantisation and Constrained Dynamics, Leuven Notes in Mathematical and Theoretical Physics, Vol. 4 , series B
- [9] P.H. Damgaard and F. De Jonghe, preprint CERN–TH-6800/93, KUL-TF-93/04, hep-th 9302050
- [10] I.A. Batalin, P.M. Lavrov and I.V. Tyutin, J. Math.Phys. **31** (1990) 1487
- [11] W. Troost, P. van Nieuwenhuizen and A. Van Proeyen, Nucl. Phys. **B333** (1990) 727